


Citramox[®] LA

150 mg/ml Suspension for injection for cattle and pigs
Amoxicillin



HIGHLY FLUID
INJECTION
SUSPENSION



KARIZOO

Citramox[®] LA

150 mg/ml Suspension for injection for cattle and pigs

Amoxicillin



Qualitative and quantitative composition

Each ml contains:

Active substance:

Amoxicillin (as trihydrate)..... 150,00 mg

List of excipients

Silica, colloidal anhydrous

Sorbitan oleate

Propylene glycol dicaprylocaprate

PHARMACEUTICAL FORM

Suspension for injection

Target species

Cattle and pigs.

Withdrawal period



Cattle

Meat and offal: 18 days
Milk: 3 days / 72 h



Porcino

Meat and offal: 20 days

Pharmacological properties

Pharmacotherapeutic group: Antibacterials for systemic use, penicillins with extended spectrum.

ATCvet code: QJ01CA04

Pack size

Cardboard box with 1 vial of 100 ml and 250 ml.

Indications for use, specifying the target species



Cattle:

Treatment of respiratory infections caused by *Mannheimia haemolytica* and *Pasteurella multocida* susceptible to amoxicillin.



Pigs:

Treatment of respiratory infections caused by *Pasteurella multocida* susceptible to amoxicillin.

Method and route of administration

15 mg amoxicillin per kg bodyweight; corresponding to 1 ml of the veterinary medicinal product per 10 kg.

Administration should be repeated after 48 hours.

Intramuscular use.

Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 28 months.

Shelf life after first opening the immediate packaging: 28 days

Special precautions for storage

Do not store above 30°C.

CITRAMOX[®] L.A. 150 mg/ml suspension for injection for cattle and pigs **Active substance:** Amoxicillin (as trihydrate) 150.00 mg. **Pharmaceutical form** Suspension for injection White to almost white suspension. **Target species:** Cattle and pigs. **Indications for use, specifying the target species** In cattle: Treatment of respiratory infections caused by *Mannheimia haemolytica* and *Pasteurella multocida* susceptible to amoxicillin. In pigs: Treatment of respiratory infections caused by *Pasteurella multocida* susceptible to amoxicillin. **Contraindications** Do not use in cases of known hypersensitivity to penicillins, cephalosporins or to any of the excipients. Do not use in cases of severe renal dysfunction with anuria and oliguria. Do not use in rabbits, hares, hamsters, guinea pigs or other small herbivores. Do not administer to Equidae, because amoxicillin – like all aminopenicillins – may adversely affect the bacterial flora of the caecum. Do not administer intravenously. **Amounts to be administered and administration route** Intramuscular use. To ensure a correct dosage and to avoid underdosing, body weight should be determined as accurately as possible. 15 mg amoxicillin per kg bodyweight; corresponding to 1 ml of the veterinary medicinal product per 10 kg. Administration should be repeated after 48 hours. **Overdose (symptoms, emergency procedures, antidotes), if necessary** Amoxicillin has a wide safety margin. In case of overdose, treatment is symptomatic. **Withdrawal period** Cattle: Meat and offal: 18 days. Milk: 3 days. Pigs: Meat and offal: 20 days. **Pharmacological properties** Pharmacotherapeutic group: Antibacterials for systemic use, penicillins with extended spectrum, ATCvet code: QJ01CA04. **Major incompatibilities** In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products. **Shelf life** Shelf-life of the veterinary medicinal product as packaged for sale: 28 months. Shelf life after first opening the immediate packaging: 28 days. **Special precautions for storage** Do not store above 30°C. **Pack size:** Cardboard box with 1 vial of 100 ml and 250 ml. **Marketing authorisation holder:** LABORATORIOS KARIZOO, S.A. Pol. Ind. La Borda, Mas Pujades 11-12 08140 Caldes de Montbui - Barcelona Spain **marketing authorisation number** 3943 ESP. **For animal treatment only. To be supplied only on veterinary prescription.**